

DATA ASSIMILATION OF GNSS ZTD

Operational and reanalyses

Martin Ridal

Magnus Lindskog (SMHI), Per Uden (SMHI), Erik Gregow (FMI)

An operational cooperation between Sweden, Norway and Finland.
Estonia, Latvia and Lithuania will join the cooperation by 2022.


- Domain is 900x960 points
- 2.5km grid spacing
- linear grid
- 65 vertical levels.
- 3 hour cycling



- MEPS currently consists of 1+9 members
- Observations used are (3Dvar):
 - Conventional observations
 - Satellite radiances (AMSU-A, MHS, IASI)
 - Scatterometer
 - Radar reflectivity)
 - GNSS ZTD
- EDA derived structure functions
- SLAF (Scaled Lagged Average Forecast) as boundaries
- From February:
 - CMEPS – 4-6 members every hour, i.e. up to 30 members
 - ECMWF-ENS at the boundaries
- Observations to come
 - MODE-S
 - GNSS slant delays

Available NGA1 at E-GVAP

Operational since June 2018
Still indicated as "test" at egvap.dmi.dk




Operational use in MetCoOp

Included are:

NGA1
METO
ROBH


Hourly data

Stations selected through a
white list



MEPS-mbr000: First Guess Departure + Bias Correction Map
db=ccma, DTG=2019-11-22 00 UTC, obname=synop, varname=apd

Observation – first guess





Variational Bias Correction

Linear predictor model:

$$b(x, \beta) = \sum_{i=0}^{N_p} \beta_i p_i(x)$$


Modified cost function:

$$J(x, \beta) = \frac{1}{2}(x - x^B)^T B^{-1}(x - x^B) + \frac{1}{2}(\beta - \beta^B)^T B_\beta^{-1}(\beta - \beta^B) + \frac{1}{2}(Hx + b(x, \beta) - y)^T R^{-1}(Hx + b(x, \beta) - y)$$



Impact of different VARBC predictor choices – 3 hour cycling


Bias and RMSE for +12,24 hour relative humidity forecasts



Small impact of introducing one more predictor in GNSS VARBC.

MEPS-mbr000: First Guess Departure Map
db=ccma, DTG=2019-11-22 00 UTC, obname=synop, varname=apd

Observation – first guess




- 3 hourly cycling
- 3 hour assimilation window, 1.5 hour before analysis time and 1.5 after
- Sub hourly GNSS ZTD data need some work
- GNSS ZTD introduced successfully (technically) every 30 minutes
- VARBC does not work properly
- To be continued...

- Same domain/resolution/levels as the operational runs
 - Updates every hour with 15 minutes cut off time
 - Up to 9 hour forecasts with output every 15 minutes
 - Rapid refresh, i.e. the first guess for upper air is operational forecast (3-6 hours)
 - Surface is cycled
 - Same observations as operational runs EXCEPT GNSS ZTD and ASCAT
-
- Sub-hourly GNSS data is used
 - Observations from XX:45 from the previous hour is used
 - Stations are selected through a white list
 - Only Swedish stations

MetCoOp nowcasting (still in development)

MNWC-preop: Observation Usage
db=ecma, DTG=2019-11-18 12 UTC, obname=synop, varname=apd



Regional Reanalysis over Europe

Time line of service and system details


2017

2018

2019

2020

2021



- 11 km (565x565 grid points), 65 levels
- Surface downscaling analysis 5.5 km (MESCAN-SURFEX)
- Start January 1961, end July 2019

Time line of service and system details



2017

2018

2019


2020

2021



- 11 km (565x565 grid points), 65 levels
- Surface downscaling analysis 5.5 km (MESCAN-SURFEX)
- Start January 1961, end July 2019


Regional Reanalysis over Europe




Regional Reanalysis over Europe

C3S_322_Lot1 Copernicus Climate Change Service - Regional Reanalysis for Europe

- 5.5 km (1080x1080 grid points)
- 106 vertical levels
- MESCAN surface analysis 5.5 km
- Continue the RRA from the FP7 UERRA project in near real time
- Deliver a RRA for Europe with an enhanced system from the early 80's (available via CDS)
- A small ensemble with 10 members at lower resolution
- A comprehensive set of output parameters, ECVs, upper air, surface and soil
- Collaboration and coordination with other reanalysis activities (e.g. Lot2 and ERA5)
- User guidance and support




Global Reanalysis → Regional Reanalysis → Surface Reanalysis




Regional Reanalysis over Europe

- GNSS ZTD will be included
- Re-processed data from the EPN EUREF Permanent Network will be used
- Data covers the period 1997–2014.
- Operational data 2015–
- 94 station in a white list



Verification of relative humidity July 2017



GBGNSS + AMSU

GBGNSS passive + AMSU

GNSS ZTD from the NGAA processing centre is used in the MetCoOp operational data assimilation.

Wish/plan:

Increase the station density

Introduce slant delays

Introduce GNSS ZTD in HARMONIE 4D-Var data assimilation.

Wish/plan:

Some code cleaning

Get VARBC in shape

Sub-hourly GNSS ZTD from Swedish stations is used in the preoperational MetCoOp nowcasting.

Wish/plan:

Get access to subhourly ZTD from all Nordic countries

Reprocessed GNSS ZTD is used in the Copernicus regional re-analysis over Europe

Wish/plan:

Everything is perfect!